# 6 SEM TDC MTMH (CBCS) C 14

2025

(May)

#### **MATHEMATICS**

(Core)

Paper: C-14

# ( Ring Theory and Linear Algebra—II )

Full Marks: 80

Pass Marks: 32

Time: 3 hours

The figures in the margin indicate full marks for the questions

- 1. (a) If F is commutative, then write the condition such that F[x] is invertible.
  - (b) Prove that every Euclidean domain possesses unity.
  - (c) Show that  $x^2 + 3x + 2$  has four zeros in  $Z_6$ .

	(d)	Let $F$ be a field. Then prove that the ring of polynomial $F[x]$ is principal ideal domain (PID).	4
	(e)	a field has at most n zeros, counting multiplicity.	6
		Or	
	7 71	Let $F$ be a field and let $f(x)$ , $g(x) \in F[x]$ with $g(x) \neq 0$ . Then prove that there exist unique polynomials $q(x)$ and $r(x)$ in $F[x]$ such that $f(x) = g(x)q(x) + r(x)$ and either $r(x) = 0$ or deg $r(x) < \deg g(x)$ .	
		A Continue of the continue of	
2.	(a)	What is the inverse of $1+\sqrt{2}$ in $Z[\sqrt{2}]$ ?	1
	(b)	Define Euclidean domain.	1
	(c)	Test the irreducibility of the polynomial $x^5 + 9x^4 + 12x^2 + 6$ in $Q$ .	2
	(d)	Prove that in a principal ideal domain, an element is irreducible if and only if it is a prime.	5
	(e)	Define unique factorization domain and prove that every field is unique factorization domain. 1+5	5=6
25/	961	( Continue	d)

Prove that  $Z[\sqrt{3}] = \{a + b\sqrt{3} \mid a, b \in Z\}$  is a Euclidean domain.

6

3. (a) Write when two linear functionals are said to be equal on a vector space V(F).

1

(b) Define invariant subspace.

1

(c) If  $S_1$  and  $S_2$  are two subsets of a vector space V(F) such that  $S_1 \subseteq S_2$ , then show that  $S_2^{\circ} \subseteq S_1^{\circ}$ .

2

(d) Prove that the subspace spanned by two subspaces each of which is invariant under some linear operator *T*, is itself invariant under *T*.

3

(e) Let V be an n-dimensional vector space over the field F and let

$$\beta = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$$

be a basis for V. Then prove that there is a uniquely determined basis

$$\beta' = \{f_1, f_2, \dots, f_n\}$$

for V' such that  $f_i(\alpha_j) = \delta_{ij}$ .

6

Let V be finite dimensional vector space over the field F and let W be a subspace of V. Then prove that

 $\dim W + \dim W^{\circ} = \dim V$ 

- 4. (a) Write about the eigenvalues and eigenvectors of the identity matrix.
  - (b) If V is n-dimensional vector space, then what is the condition that the linear operator T is diagonalizable?
  - (c) Test the diagonalizability of the following matrix:

$$\begin{bmatrix} \frac{1}{2} & \frac{3}{2} \\ \frac{3}{2} & \frac{1}{2} \end{bmatrix}$$

(d) Define minimal polynomial and show that the minimal polynomial of the real matrix

$$\begin{bmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{bmatrix}$$

is 
$$(x-1)(x-2)$$
.

1

1

If f(x) be the characteristic polynomial of T, then prove that  $f(T) = \hat{0}$ .

any times product space

6

5. (a) Write the only vector that is orthogonal to itself.

(b) Define orthogonal complement.

1

1

(c) If  $\alpha$ ,  $\beta$  are vectors in an inner product space V, then prove that

$$\|\alpha + \beta\| \le \|\alpha\| + \|\beta\|$$

Or 40 letter well

If  $W_1$  and  $W_2$  are subspaces of a finite dimensional inner product space, then prove that

$$(W_1 + W_2)^{\perp} = W_1^{\perp} \cap W_2^{\perp}$$

(d) If  $\beta = \{\alpha_1, \alpha_2, \dots, \alpha_m\}$  is any finite orthonormal set in an inner product space V, and if  $\beta$  is any vector in V, then prove that

$$\sum_{i=1}^{m} \left| (\beta, \alpha_i) \right|^2 \le \left\| \beta \right\|^2$$

In an inner product space, prove that

### $|(\alpha, \beta)| \le ||\alpha|| ||\beta||$

- 6. (a) Write the two self-adjoint operators on any inner product space V(F).
  - (b) Define normal operator. 1
  - (c) If  $T_1$  and  $T_2$  are normal operators on an inner product space with the property that either commutes with the adjoint of the other, then prove that  $T_1$   $T_2$  is also normal operator.
  - (d) Let V be the direct sum of its subspaces  $W_1$  and  $W_2$ . If  $E_1$  is the projection on  $W_1$  along  $W_2$ , and  $E_2$  is the projection on  $W_2$  along  $W_1$ , then prove that—
    - (i)  $E_1 + E_2 = I$ ;
    - (ii)  $E_1 E_2 = \hat{0}, E_2 E_1 = \hat{0}.$
  - (e) If  $T_1$  and  $T_2$  are self-adjoint linear operators on an inner product space V, then prove that (i)  $T_1 + T_2$  is self-adjoint and (ii) if  $T_1 \neq \hat{0}$  and a is a non-zero scalar, then  $aT_1$  is self-adjoint iff a is real.

5

1

2

Apply the Gram-Schmidt process to the vectors (1,0,0), (1,1,0), (1,1,1) to obtain an orthonormal basis for  $V_3(R)$  with the standard inner product.

\*\*\*