

Total No. of Printed Pages—8

**3 SEM TDC MTH M 2**

**2 0 1 7**

( November )

**MATHEMATICS**

( Major )

Course : 302

**( Coordinate Geometry and Algebra—I )**

*Full Marks : 80*

*Pass Marks : 32/24*

*Time : 3 hours*

*The figures in the margin indicate full marks  
for the questions*

**GROUP—A**

**( Coordinate Geometry )**

**SECTION—I**

**( 2-Dimension )**

( Marks : 27 )

1. (a) What will be the equation of the circle  $(x - h)^2 + (y - k)^2 = r^2$ , when the origin is transferred to the point  $(h, k)$ ? 1

- (b) Prove that if  $ax^2 + 2hxy + by^2 = 1$  and  $a'x'^2 + 2h'xy + b'y^2 = 1$  represent the same conic and the axes are rectangular, then show that

$$(a - b)^2 + 4h^2 = (a' - b')^2 + 4h'^2 \quad 2$$

- (c) The axes are rotated through an angle  $60^\circ$  without changing the origin. If the coordinates of a point are  $\left(\frac{1}{2}, \frac{5\sqrt{3}}{2}\right)$  in old system, what would be its coordinates in new system? 2

2. (a) State the name of the geometrical figure represented by the equation  $xy = 0$ . 1
- (b) Find the equations of the straight lines which pass through the origin and whose distance from  $(h, k)$  are equal to  $d$ . 2

- (c) Prove that the lines represented by

$$ax^2 + 2hxy + by^2 + \lambda(x^2 + y^2) = 0$$

have the same pair of bisectors for all values of  $\lambda$ . Interpret the case for  $\lambda = -(a + b)$ .

2+2=4

Or

If the straight lines represented by the equation

$$x^2(\tan^2 \phi + \cos^2 \phi) - 2xy \tan \phi + y^2 \sin^2 \phi = 0$$

make angles  $\alpha$  and  $\beta$  with the  $x$ -axis, show that  $\tan \alpha - \tan \beta = 2$ . 4

- (d) Find the condition that one of the lines given by  $ax^2 + 2hxy + by^2 = 0$  may be perpendicular to one of the lines given by  $a'x^2 + 2h'xy + b'y^2 = 0$ . 5

Or

Prove that the equation

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$

represents a pair of parallel straight lines if  $\frac{a}{h} = \frac{h}{b} = \frac{g}{f}$ .

3. (a) State True or False : 1

When the focus lies on the directrix, the conic section is a pair of lines.

- (b) Find the equation of the polar of the point (2, 3) with respect to the conic

$$x^2 + 3xy + 4y^2 - 5x + 3 = 0 \quad 2$$

- (c) Find the condition that the pair of lines  $Ax^2 + 2Hxy + By^2 = 0$  may be conjugate diameter of the conic

$$ax^2 + 2hxy + by^2 = 1 \quad 3$$

Or

Find the equation of the diameter of the conic  $15x^2 - 20xy + 16y^2 = 1$  conjugate to the diameter  $y + 2x = 0$ .

- (d) Find the equation of the chord of contact of tangents from a given point  $(x_1, y_1)$  to the conic

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0 \quad 4$$

Or

Reduce the equation

$$9x^2 - 24xy + 16y^2 - 18x - 101y + 19 = 0$$

to the standard form.

SECTION—II

( 3-Dimension )

( Marks : 18 )

4. (a) If  $O$  is origin and  $P(a, b, c)$ , then write the direction cosines of the line  $OP$ . 1

- (b) A plane cuts the axes at  $A, B, C$  and the centroid of the triangle is  $(a, b, c)$ . Find the equation of the plane. 3

Or

Find the equation of the plane passing through the point  $(3, -2, 6)$  and through the  $x$ -axis.

- (c) Find the equations of the line passing through the point  $(2, 1, 0)$  and parallel to the line joining the points  $(1, 5, 2)$  and  $(3, 0, -1)$ . 2

- (d) Put the equations

$$4x - 4y - z + 11 = 0 = x + 2y - z - 1$$

of a line in the symmetrical form. 4

Or

Find the distance of the point  $(-3, 1, 1)$  from the plane  $2x + y - 4z + 6 = 0$  measured parallel to the line

$$\frac{x}{2} = \frac{y}{-2} = \frac{z}{1}$$

5. (a) Fill in the blank : 1

The shortest distance between the two lines is their common \_\_\_\_.

- (b) Find the shortest distance between the line

$$ax + by + cz + d = 0 = a'x + b'y + c'z + d'$$

and the z-axis. 3

- (c) Show that the shortest distance between the lines  $x - y + z = 0 = 2x - 3y + 4z$  and  $x + y + 2z - 3 = 0 = 2x + 3y + 3z - 4$  is  $\frac{13}{\sqrt{66}}$ . 4

Or

Find the length and the equations of the shortest distance between the lines

$$\frac{x-1}{2} = \frac{y+8}{-7} = \frac{z-4}{5} \text{ and } \frac{x-1}{2} = \frac{y-2}{1} = \frac{z-6}{-3}$$

GROUP—B

( Algebra—I )

( Marks : 35 )

6. (a) State True or False : 1  
Addition of natural number in binary composition is not associative.
- (b) Define a quaternion group. 1
- (c) Show that a finite semigroup in which cross-cancellation holds is an Abelian group. 2

(d) Answer any two : 3×2=6

(i) Show that a subgroup of a cyclic group is cyclic.

(ii) Let  $H, K$  be subgroup of  $G$ . Show that  $HK$  is a subgroup of  $G$  if and only if  $HK = KH$ .

(iii) Show that for elements  $a, b$  in a group  $G$ , the equations  $ax = b$  and  $ya = b$  have unique solutions for  $x$  and  $y$  in  $G$ .

7. Answer any two : 5×2=10

(a) If a group has finite number of subgroups, then show that it is a finite group.

(b) If  $G$  be a group and  $a, b \in G$ , such that (i)  $ab = ba$  and (ii)  $O(a), O(b) = 1$ , then show that  $O(ab) = O(a)O(b)$ .

(c) Prove that a nonempty subset  $H$  of a group  $G$  is a subgroup of  $G$ , iff—

(i)  $a, b \in H \Rightarrow ab \in H$ ;

(ii)  $a \in H, a^{-1} \in H$ .

8. (a) Define a simple group. 1

(b) Prove that every quotient group of a cyclic group is cyclic. 4

Or

If  $G$  is a group such that  $\frac{G}{Z(G)}$  is cyclic, where

$Z(G)$  is centre of  $G$ , then show that  $G$  is Abelian.

5

(c) Answer any two : 5×2=10

(i) Show that a subgroup  $H$  of a group  $G$  is normal in  $G$  if and only if

$$g^{-1}hg \in H \quad \forall h \in H, g \in G.$$

(ii) If  $H$  and  $K$  be two subgroups of a group  $G$ , where  $H$  is normal in  $G$ , then prove that

$$\frac{HK}{H} \cong \frac{K}{H \cap K}$$

(iii) Find the regular permutation groups isomorphic to the multiplicative group  $G = \{1, -1, i, -i\}$ .

\*\*\*